

# Hard X-ray Variability in M82: Evidence for a Nascent AGN?

A. Ptak<sup>1</sup> & R. Griffiths

Carnegie Mellon University, Dept. of Physics, Pittsburgh, PA 15213

## ABSTRACT

We report on the detection of hard (2-10 keV) X-ray variability in the starburst galaxy M82 over the course of 9 *ASCA* observations. Long-term variability occurred on a time scale of days, with a change in flux by a factor of up to  $\sim 4$ , corresponding to a point-source luminosity of  $L_{2-10 \text{ keV}} \sim 6 \times 10^{40} \text{ ergs s}^{-1}$ . Short-term variability with an amplitude of  $\sim 1.4$  on a timescale of hours was observed during the longest observation. This demonstrates that a large fraction of the hard X-ray emission of M82 (depending on the flux state) is from a compact region and is probably due to an accreting source. The 2-10 keV luminosity of the source is a lower limit to its Eddington luminosity, implying a blackhole mass of at least  $\sim 460 M_{\odot}$ , or a mass intermediate to that of normal AGN and stellar-mass blackhole candidates.

*Subject headings:* galaxies: starburst – galaxies: individual (M82) – X-rays: galaxies

## 1. Introduction

Hard X-rays (2 – 10 keV) were marginally detected from M82 by *Uhuru* (Forman et al. 1978), *Ariel-5* (Cooke et al. 1978; McHardy et al. 1981), and *HEAO-1* (Piccinotti et al. 1982), making it one of the earliest extragalactic hard X-ray sources, accurately positioned using the HEAO-A3 instrument (Griffiths et al. 1979). These detections were confirmed by the *Einstein MPC* (Watson, Stanger, & Griffiths 1984; Fabbiano 1988), *EXOSAT ME* (Schaaf et al. 1989) and *Ginga* (Ohashi et al. 1990). Schaaf et al. speculated that the hard X-ray flux from M82 was due to inverse-Compton scattering of infrared photons off relativistic electrons (originally suggested by Hargrave [1974]). Ohashi et al. claimed that a bremsstrahlung fit was preferred over a non-thermal power-law for the 2-20 keV *Ginga* spectrum, suggesting a thermal origin to the flux. However, the lack of a strong (equivalent width  $\sim 1 \text{ keV}$ ) Fe-K line detection is problematic for a thermal model, unless the abundances in the nucleus of M82 are significantly subsolar (which would not be expected to be the case for a starburst galaxy). A short *BBXRT* observation found that the 0.5-10.0 keV spectrum of M82 consists of at least two components, with the hard component described well by either a power-law model or a thermal plasma model (Petre 1993). M82 was originally observed by *ASCA* in 1993 during the Performance Verification (PV) phase of the mission. These data confirmed the *BBXRT* results and were analyzed by several groups (Moran & Lehnert 1997; Ptak et al. 1997; Tsuru et al. 1997).

Moran & Lehnert came to the conclusion that the hard flux originated in IC scattering of infrared flux based on the good correlation of *ROSAT* “hardness” and infrared images. Ptak et al. speculated that a variable X-ray binary detected by *Einstein* (Watson, Stanger & Griffiths 1984) and *ROSAT* (Collura et al. 1994) *could* account for a large fraction of the 2-10 keV flux from M82 *if* its X-ray spectrum was similar to the spectrum of the hard component in M82. Similarly, Tsuru et al. suggested that M82 has varied significantly in the 2-10 keV bandpass,

---

<sup>1</sup>email: ptak@astro.phys.cmu.edu

and accordingly the hard X-ray emission is most likely due to an accreting binary or AGN. However, all fluxes prior to *ASCA* were from *non-imaging* detectors, and accordingly any flux comparison is inherently suspect (i.e., due to the collimation of diffuse flux and spurious sources in the non-imaging detector’s field of view). For example, M81, which is a variable 2-10 keV source, lies  $\sim 40'$  from M82. Here we report on 2-10 keV intensity and spectral variability detected in M82 from a series of monitoring observations by *ASCA*.

## 2. The *ASCA* Data

The *ASCA* observations were performed during the period Mar. 6, 1996 through Nov. 11, 1996 at 9 intervals (see Table 1). Briefly, *ASCA* (Tanaka et al. 1994) comprises two solid-state imaging spectrometers (SIS; hereafter S0 and S1) with an approximate bandpass of 0.4-10.0 keV and two gas imaging spectrometers (GIS; hereafter G2 and G3) with an approximate bandpass of 0.8-10.0 keV. The SIS observations were done in 1-ccd mode<sup>2</sup>, and due to calibration problems below 0.6 keV we only consider SIS data in the 0.6-10.0 keV bandpass (most of our conclusions are based on data in the 2.0-10.0 keV bandpass). Because the 1993 observation showed that the 2-10 keV emission from M82 is unresolved (Tsuru et al. 1997, Ptak et al. 1999), we chose source regions appropriate for a point source, namely  $6'$  for the GIS detectors and  $4'$  for the SIS detectors. The background was determined from the remaining counts in the CCDs beyond  $5'$  for the SIS, and from a  $7$ - $12'$  annulus for the GIS data. The spectra from each set of detectors were combined to yield a SIS and GIS spectrum for each observation, although we also performed fits to each spectrum separately to check for anomalies. Prior to spectral fitting the data were binned to a minimum of 20 counts per bin to allow the use of the  $\chi^2$  statistic.

## 3. Results

### 3.1. Long-term Variability

A good fit to the M82 *ASCA* PV spectra required at least two components: a thermal plasma model dominating the soft flux below 2 keV and a power-law or thermal bremsstrahlung model dominating above 3 keV (Tsuru et al. 1997, Ptak et al. 1997, Moran & Lehnert 1997). Accordingly, we fit the two spectra from each observation with a thermal plasma plus a power-law model, with only the power-law (dominating above 2-3 keV) results being of interest in the present work. Note that our motivation for fitting the spectra here is for the purpose of determining fluxes rather than determining the precise values of spectral parameters. These fits were acceptable ( $\chi^2_\nu = 0.95 - 1.10$ ). Figure 1 shows the 2-10 keV long-term lightcurve based on (observed) fluxes inferred from these fits. We also fit the data in the 3-10 keV bandpass only with a simple power-law and found similar values for both fluxes and the power-law slope, showing that the soft component is not adversely affecting our results (in both sets of fits the mean column density absorbing the power-law was on the order of  $2 \times 10^{22} \text{ cm}^{-2}$ , similar to the PV observation value of  $1.9 \pm 1.0 \times 10^{22} \text{ cm}^{-2}$  given in Ptak et al. 1997). Long-term variability is evident, with a change in 2-10 keV flux from  $\sim 1.4 \times 10^{-11} \text{ ergs cm}^{-2} \text{ s}^{-1}$  to  $\sim 5.2 \times 10^{-11} \text{ ergs cm}^{-2} \text{ s}^{-1}$ . The soft flux from M82 is known to be extended, with some contribution from point sources that are variable (c.f., Collura et al. 1994, Bregman

---

<sup>2</sup>see [http://adfwww.gsfc.nasa.gov/asca/processing\\_doc/GS/intro.html#highlights](http://adfwww.gsfc.nasa.gov/asca/processing_doc/GS/intro.html#highlights)

Table 1. Observation Log

| ID | Date     | Duration<br>(ks) | Exposure<br>(ks) | Count Rates <sup>a</sup><br>(cts/s) | Angle <sup>b</sup><br>(arcmin) |
|----|----------|------------------|------------------|-------------------------------------|--------------------------------|
| 1  | 3/23/96  | 20.0             | 12.9-13.6        | 0.89,0.73,0.47,0.60                 | 5.7,7.8,7.5,5.1                |
| 2  | 4/15/96  | 18.2             | 6.9-8.7          | 1.25,1.07,0.75,0.95                 | 5.8,8.2,7.6,5.3                |
| 3  | 4/21/96  | 24.1             | 12.4-12.9        | 1.15,0.94,0.66,0.84                 | 5.8,8.1,7.6,5.2                |
| 4  | 4/24/96  | 64.3             | 29.3-33.3        | 1.15,0.92,0.66,0.82                 | 5.8,8.3,7.7,5.5                |
| 5  | 5/13/96  | 18.6             | 8.1-9.4          | 0.78,0.64,0.41,0.51                 | 6.2,8.6,8.0,5.7                |
| 6  | 5/5/96   | 19.3             | 11.5-14.4        | 0.91,0.76,0.49,0.60                 | 6.2,8.6,8.0,5.7                |
| 7  | 10/14/96 | 25.5             | 7.6-8.6          | 0.59,0.51,0.30,0.36                 | 6.5,8.9,8.4,5.9                |
| 8  | 11/14/96 | 20.4             | 12.0-12.5        | 0.68,0.56,0.33,0.43                 | 6.1,8.2,7.9,5.5                |
| 9  | 11/26/96 | 14.0             | 7.6-9.8          | 0.76,0.67,0.43,0.56                 | 5.9,7.7,7.6,4.7                |

Note. — The count rates cited are for the full bandpass of each detector ( $\sim 0.6 - 10.0$  keV and  $\sim 0.8 - 10.0$  keV for the SIS and GIS, respectively). The signal-to-noise ratio of the observations varied from 45-177, with a mean of 88.

<sup>a</sup>Background-subtracted count rates for S0, S1, G2, G3.

<sup>b</sup>Off-axis angle for S0, S1, G2, G3.




Fig. 1.— top) The 0.5-2.0 keV (filled stars) and 2-10 keV (filled circles) long-term light curve from M82. The fluxes are not corrected for extinction and are inferred from a two-component fit to the integrated spectrum from each observation. The error bars are at a level of 10%, which is a reasonable estimate of the relative uncertainty in flux from observation to observation. (bottom) The best-fitting power-law energy indices resulting from the fits used to determine the fluxes. The error bars show the 90% confidence intervals (see text). The power-law dominates the 2-10 keV bandpass.

Schullman & Tomisaka 1995). The 0.5-2.0 keV fluxes derived from the same fits (also shown in Figure 1) remained within  $\sim 15\%$  of the mean, demonstrating that the variability is occurring predominately above 2 keV. The stability of the soft flux also indicates that the observed variability is not due to instrumental effects (which would likely affect the entire bandpass in a similar fashion) or the explosion of a supernova which would likely result in a softer spectrum. The mean 0.5-2.0 keV flux in these observations is  $1.1 \times 10^{-11}$  ergs  $\text{cm}^{-2}$   $\text{s}^{-1}$ , comparable to the value cited in Ptak et al. (1997) for the PV observation ( $9.4 \times 10^{-12}$  ergs  $\text{cm}^{-2}$   $\text{s}^{-1}$ ; the 2-10 keV PV flux is consistent with the flux from the 8th observation given here). Note that some of the variability below 2 keV may be due to the same source(s) causing the hard variability, but the precise quantification of this is uncertain since the level of variability is on the order of systematic errors. The power-law energy index resulting from these fits is shown in the bottom panel of Figure 1. The errors bars shown are the 90% confidence based on  $\Delta\chi^2 = 17$  (11 interesting parameters), with similar results being obtained for  $\Delta\chi^2 = 4.6$  (2 interesting parameters) in the 3-10 keV fits. The typical 90% confidence on the energy index was  $\sim 0.3$ . In most of the observations, the energy index was consistent with the mean value of 1.0, however, the energy index in the first observation was significantly steeper,  $\alpha = 1.68$  (1.42-1.97) (the PV

observation power-law slope was  $0.76^{+0.22}_{-0.21}$ ).




Fig. 2.— The resultant spectrum when the lowest flux spectrum is subtracted from the highest flux spectrum. The GIS data are marked with filled circles, the SIS data are unmarked. A power-law fit to the data is shown with a solid line (SIS data) and a dotted line (GIS data). The ratio of the data to the model is shown in the bottom panel.

We also investigated the residual spectrum when the lowest flux state spectrum is subtracted as background from the other spectra. The motivation for this was to isolate the spectrum of the variable source by subtracting the (extended) soft flux, as well as any other sources of hard X-ray flux (thermal emission, IC scattered photons, other point sources, and background). We subtracted the spectra prior to any binning (or background subtraction). Figure 2 shows the results of a power-law fit to the highest flux spectrum residuals. The power-law fit is excellent in the 2-10 keV bandpass, as expected, but interestingly a soft “tail” is also present, at a flux level (in this case) of  $\sim 1 \times 10^{-12}$  ergs cm $^{-2}$  s $^{-1}$ , or about 10% of the total 0.5-2.0 keV flux from M82, which is typical of the scatter in the 0.5-2.0 keV lightcurve shown in Figure 1. A detailed discussion of these fits is beyond the scope of this paper, but we note that the soft excess must be treated cautiously since a 10% variation could be a systematic effect resulting from the subtraction of separate observations (although note this component appears consistently among the SIS and GIS data suggesting that this is not the case).

Since the pointing accuracy of *ROSAT* is significantly better than that of *ASCA*, and the 0.5-2.0 keV flux observed by both satellites must have a common origin, the position of the hard X-ray flux can best be estimated by comparing the centroids of the hard and soft X-ray emission. We found that the centroids of the 0.5-2.0 keV and 3.0-10.0 keV flux are coincident to within  $\sim 10''$  ( $\sim 200$  pc), indicating that the emission is nuclear, i.e., the 0.5-2.0 keV emission

is known to originate in the nucleus from *Einstein* and *ROSAT* HRI observations (Watson, Stanger & Griffiths 1984; Bregman et al. 1995). We investigated the radial profile distribution of the lowest flux state observation (ID 7 in Table 1) and of the most variable observation (ID 4, see below) and found that the 3-10 keV emission in both cases is unresolved. A radial Gaussian fit to the SIS0 radial profiles (see Ptak 1997, Ptak et al. 1999 for the method) yielded an upper limit to the half-light radius of the emission on the order of  $30''$  ( $\sim 0.5$  kpc). These extents are comparable to the spatial extent found in Ptak (1997) and Ptak et al. (1999) for the *ASCA* PV M82 SIS data. This result indicates that the source of variability is most likely a single nuclear source rather than several X-ray binaries distributed over M82 (although a small *nuclear* population of binaries cannot be excluded as a possibility).

### 3.2. Short-Term Variability




Fig. 3.— The short-term, 2-10 keV SIS (dashed lines marked with stars) and GIS lightcurve (solid lines marked with filled circles) resulting from each individual lightcurve concatenated onto a single plot. The duration of each observation (delimited by vertical dotted lines) is 25 hours, and X-axis labels give the start times of the observations. The background rate is also shown in the figure (at a mean rate of  $0.031 \text{ counts s}^{-1}$ ,  $\sim 5\%$  of the total, and  $0.044 \text{ counts s}^{-1}$ ,  $\sim 7\%$  of the total, for the SIS and GIS, respectively.). Note that given the *ASCA* PSF, several percent of the source flux is scattered into the background regions used. The size of each bin was  $\sim 4000$  s.

Figure 3 shows the short-term lightcurves of each observation appended into a single plot. The lightcurve from the 4th observation is shown in Figure 4, where a large variation is evident.

The source varied by  $\sim 20\%$  from the mean flux of the observation, or a peak-to-valley variation of  $\sim 1.4$ . This type of fluctuation was observed so clearly in this observation only, with hints of variability (but typically at a lower level) in the remaining observations. A sine function fit to the lightcurve in Figure 4 results in a period of  $\sim 15$  hours. We “detrended” the long-term light-curve by subtracting the mean count rate of each individual observation, and found that trend observed in Figure 4 does not extend to the other lightcurves, i.e., this “QPO” is not consistent with an underlying periodicity superimposed on long-term variations. An alternative possibility is that this is a flare with a duration of  $\sim 15$  hours that caused an increase in the flux by  $\sim 40\%$ . Note, however, that the durations of the other observations were typically only 5 hours, so strong conclusions cannot be drawn concerning the nature of the short-term variability until a longer, contiguous observation of M82 is made in the 2-10 keV bandpass.




Fig. 4.— The short-term, 2-10 keV SIS and GIS lightcurves (marked as in Figure 3) from the 4th observation (the observation with the longest duration) is shown. The curve shows a sinusoidal fit to the GIS data (similar results are obtained from the SIS data), with a best-fit period of 14.7 hours ( $\chi^2/\text{dof} = 100.2/50$ ).

#### 4. Discussion

We have presented data showing that a variable source of 2-10 keV flux is present in M82, located within  $\sim 10''$  (170 pc) of the nucleus. This source may be associated with the variable X-ray source discussed in Collura et al., although higher angular resolution is necessary to assess

this possibility (with *ASCA* we can only note that the hard and soft flux are coincident to within  $\sim 10''$ , which the Collura et al. source, which can dominate the soft flux, is within  $\sim 5''$  of the nucleus of M82). Here for the first time we show that a variable, compact source is not only producing hard emission, but can also *dominate* the hard flux from M82. There is a suggestion of periodic or semi-periodic variability, similar to that observed in some galactic X-ray binaries (c.f., Verbunt 1993).

The marginal detection of Fe-K $\alpha$  emission at  $\sim 6.7$  keV in the 1993 *ASCA* observation (with M82 at a 2-10 keV flux of  $\sim 2.0 \times 10^{-11}$  ergs cm $^{-2}$  s $^{-1}$ ) implies that at least some of the hard component is thermal (Ptak et al. 1997, see also the *BeppoSAX* Fe-K detection given in Cappi et al. 1999a). The source also appears to exhibit spectral variability, although predominately just between the first and remaining observations. No correlation between spectral slope and flux is observed, although the statistics are limited.

Assuming a distance to M82 of 3.6 Mpc (Freedman et al. 1994), the difference between the highest and lowest 2-10 keV fluxes observed corresponds to a luminosity of  $5.9 \times 10^{40}$  ergs s $^{-1}$ . For comparison, the brightest nearby blackhole candidate (BHC) is SMC X-1 with a 2-10 keV luminosity of  $6 \times 10^{38}$  ergs s $^{-1}$  (Verbunt 1993). The timescale of this variability requires the source of this hard X-ray flux to be compact, and it is most likely an accreting system with a lower-limit on the blackhole mass of  $\sim 460 M_{\odot}$  unless the high flux rate is exceeding the Eddington limit for the system, the emission is anisotropic, and/or the relativistic motions are significant. Given the high luminosity of this source (i.e., in a galaxy not known to be harboring an AGN), these caveats should not be taken lightly. As discussed in Komossa & Bade (1999), other exotic possibilities for luminous variable X-ray sources such as this include a radio supernova, the tidal disruption of a star by a massive blackhole, and a gamma ray burst. However, the lack of a monotonic decline in the lightcurve is problematic for these possibilities<sup>3</sup>. Taken at face value, this mass estimate is intermediate between the  $10^6$ – $10^9 M_{\odot}$  systems found in active galactic nuclei (AGN) and stellar-mass blackhole systems, and this source may represent either an existing low-luminosity AGN (with a mass on the order of  $10^6 M_{\odot}$  and a very low accretion rate), or a “nascent” AGN. By the latter possibility we mean a blackhole system that is gradual growing in mass to eventually become a massive AGN. The supernova rate in M82 is  $\sim 0.1$  yr $^{-1}$  (Van Buren & Greenhouse 1994), and over a typical starburst lifetime of  $10^7$  years, of order  $10^6$  solar masses of compact matter (i.e., neutron stars and blackholes) should be produced. We are therefore speculating that on the order of at least  $\sim 0.05\%$  of this matter may have coalesced in the nucleus.

It would be of interest to distinguish between the AGN and massive BHC scenarios. Note that the lightcurve of this source does not show the exponential decay characteristic of blackhole novae (c.f., Figure 3 in Tanaka & Shibazaki 1996), although a more systematic monitoring of M82 would be necessary to compare the temporal properties of this source to Galactic BHC. If the accretion disk or region in this source is optically-thick, then a  $M > 10^7 M_{\odot}$  blackhole accretion disk would emit a blackbody spectrum with a temperature less than  $10^6$  K, while in contrast accretion disk temperatures for a blackhole candidate are typically on the order of  $10^6$ – $10^7$  K (c.f., Frank et al. 1992). Therefore, the *unambiguous* detection of a soft “excess” that can be attributed to this source alone (as should be possible with the arcsecond resolution

---

<sup>3</sup> Komossa & Bade favor variability in the absorption (in either column density or ionization state) of an AGN as the explanation for the observed lightcurves of NGC 5905 and IC3599, but in our case the available of *ASCA* data above 2 keV rules out this possibility.



of *AXAF*) would yield an important clue. For example, Colbert & Mushotzky (1999) have analyzed the X-ray data available for a sample of nearby normal galaxies and found that bright, extranuclear point-sources are common. The soft spectra of these sources were fit well with disk models suggesting blackhole masses in the range of  $10^{2-4} M_{\odot}$ , consistent with the source in M82 (which may not have settled into the dynamical center of M82 yet). On the other hand, if the accretion mode is advection-dominated, then there should be no soft excess (note that in this case the relative steepness of the 2-10 keV emission implies that the accretion rate must exceed  $\sim 10^{-3}$  in Eddington units, c.f. Ptak et al. 1998). The unambiguous association between this source and counterparts in other wavebands (particularly in the radio where AGN candidates in M82 have been proposed; see Seaquist, Frayer & Frail 1997 and references therein, but also see Allen & Kronberg 1998) would go long way towards resolving this issue.

This paper made extensive use of the NASA’s High Energy Astrophysics Science Archive Research Center and Astrophysics Data System Abstract Service databases, and the NASA/IPAC Extragalactic Database. We would also like to thank the referee for useful comments.

## REFERENCES

- Allen, M. & Kronberg, P. 1998, *ApJ*, 502, 218
- Cappi, M. et al. 1999a, to appear in “Dal nano- al tera-eV: tutti i colori degli AGN”, third Italian conference on AGNs, Roma, Memorie S.A.It, in press, astro-ph/9809325.
- Cappi, M. et al. 1999b, *Advances in Space Research*, in *Proceedings of 32nd Sci. Ass. of COSPAR*, in press, astro-ph/9902021.
- Collura, A., Shulman, E., Reale, F., & Bregman, J. 1994, *ApJ*, 420, L63
- Cooke, B. et al. 1978, *MNRAS*, 182, 489
- Fabbiano, G. 1988, *ApJ*, 330, 672
- Forman, W. et al. 1978, *ApJS*, 38, 357
- Frank, J., King, A., & Raine, D. 1992, *Accretion Power in Astrophysics* (Cambridge: Cambridge University Press)
- Freedman, W. et al. 1994, *ApJ*, 427, 628
- Griffiths, R., Doxsey, R., Johnston, M., Schwartz, D., & Blades, J. 1979, *ApJ*, 230, L21
- Komossa, S. & Bade, N. 1999, *A&A*, in press, astro-ph/9901141
- Hargrave, P. 1974, *MNRAS*, 168, 491
- McHardy, I., Lawrence, A., Pye, J. & Pounds, K. 1981, *MNRAS*, 197, 893
- Moran, E. & Lehnert, M. 1997, *ApJ*, 478, 172
- Ohashi, T., Makishima, K., Tsuru, T., Takano, S., Koyama, K., & Stewart, G. 1990, *ApJ*, 368, 180
- Petre, R. 1993, in *The Nearest Active Galaxies*, eds. J. Beckman, L. Colina, and H. Netzer, p. 117
- Piccinotti, G., Mushotzky, R., Boldt, E., Holt, S., Marshall, F., Serlemitsos, P., & Shafer, R. 1982, *ApJ*, 253, 485
- Ptak, A., Serlemitsos, P., Yaqoob, T., & Mushotzky, R. 1997, *AJ*, 113, 1286
- Ptak, A. 1997, Ph.D. thesis, University of Maryland
- Ptak, A., Yaqoob, T., Mushotzky, R., Serlemitsos, P. & Griffiths, R. 1998, *ApJ*, 501, L37
- Ptak, A., Mushotzky, R., Yaqoob, T., & Serlemitsos, P. 1999, *ApJ*, submitted
- Schaaf, R., Pietsch, W., Biermann, P., Kronberg P., & Schmutzler, T. 1989, *ApJ*, 336, 722
- Seaquist, E., Frayer, D. & Frail, D. 1997, *ApJ*, 487, L131
- Tanaka, Y., Inoue, H., & Holt, S. 1994, *PASJ*, 46, L37
- Tsuru, T., Awaki, H., Koyama, K. & Ptak, A. 1997, *PASJ*, 49, 619
- Van Buren, D. & Greenhouse, M. 1994, *ApJ*, 286, 144
- Verbunt, F. 1993, *ARA&A*, 31, 93
- Watson, M., Stanger, V. & Griffiths, R. 1984, *ApJ*, 286, 144